



Mark Scheme (Results)

Summer 2024

Pearson Edexcel Advanced Subsidiary GCE
In Biology B (8BI0)
Paper 01: Core Cellular Biology and
Microbiology

Edexcel and BTEC Qualifications

Edexcel and BTEC qualifications are awarded by Pearson, the UK's largest awarding body. We provide a wide range of qualifications including academic, vocational, occupational and specific programmes for employers. For further information visit our qualifications websites at www.edexcel.com or www.btec.co.uk. Alternatively, you can get in touch with us using the details on our contact us page at www.edexcel.com/contactus.

Pearson: helping people progress, everywhere

Pearson aspires to be the world's leading learning company. Our aim is to help everyone progress in their lives through education. We believe in every kind of learning, for all kinds of people, wherever they are in the world. We've been involved in education for over 150 years, and by working across 70 countries, in 100 languages, we have built an international reputation for our commitment to high standards and raising achievement through innovation in education. Find out more about how we can help you and your students at: www.pearson.com/uk

Summer 2024

Question Paper Log Number P74474A

Publications Code 8BI0_01_2406_MS

All the material in this publication is copyright ©

Pearson Education Ltd 2024

General Marking Guidance

- All candidates must receive the same treatment. Examiners must mark the first candidate in exactly the same way as they mark the last.
- Mark schemes should be applied positively. Candidates must be rewarded for what they have shown they can do rather than penalised for omissions.
- Examiners should mark according to the mark scheme not according to their perception of where the grade boundaries may lie.
- There is no ceiling on achievement. All marks on the mark scheme should be used appropriately.
- All the marks on the mark scheme are designed to be awarded. Examiners should always award full marks if deserved, i.e. if the answer matches the mark scheme. Examiners should also be prepared to award zero marks if the candidate's response is not worthy of credit according to the mark scheme.
- Where some judgement is required, mark schemes will provide the principles by which marks will be awarded and exemplification may be limited.
- When examiners are in doubt regarding the application of the mark scheme to a candidate's response, the team leader must be consulted.
- Crossed out work should be marked UNLESS the candidate has replaced it with an alternative response.

Question Number	Answer	Additional Guidance	Mark
1(a)(i)	<p>The only correct answer is C</p> <p>A is incorrect because magnesium ions are not incorporated into amino acids or nucleotides</p> <p>B is incorrect because magnesium ions are not incorporated into amino acids</p> <p>D is incorrect because magnesium ions are not incorporated into nucleotides</p>		(1)

Question Number	Answer	Additional Guidance	Mark
1(a)(ii)	<p>An explanation that makes reference to the following:</p> <ul style="list-style-type: none"> because water is {(di)polar } (solvent) (1) ions form charged attractions / electrostatic attractions (with it) (1) 	<p>ACCEPT charged / hydrophilic</p> <p>ACCEPT form hydrogen bonds</p>	(2)

Question Number	Answer	Additional Guidance	Mark
1(b)	<p>An answer that makes reference to the following:</p> <ul style="list-style-type: none"> • more nitrate ions are taken up in the presence of oxygen (1) • nitrate ions are taken up faster in the presence of oxygen (1) • Nitrates are taken up by (diffusion and) active transport in presence of oxygen/ nitrates only taken up by diffusion without oxygen present (1) 	ACCEPT converse	(2)

Question Number	Answer	Additional Guidance	Mark
2(a)	<p>An answer that makes reference to the following:</p> <ul style="list-style-type: none"> • three identical chromosomes at position 21 (1) • two identical chromosomes at positions 20, 22 and 23 (1) 		(2)

Question Number	Answer	Additional Guidance	Mark
2(b)	<ul style="list-style-type: none"> • part of one chromosome drawn on the other chromosome (1) 		(1)

Question Number	Answer	Additional Guidance	Mark
2(c)(i)	<p>An answer that makes reference to the following:</p> <ul style="list-style-type: none"> • height in m² calculated (1) • BMI calculated (1) 	<p>(1.463 × 1.463 =) 2.140369 Accept rounding - 2.14, 2.1</p> <p>21.1 (1) (=21.1178539)</p> <p>Accept (if rounded to 2.14) 21.1214953 Accept (if rounded to 2.1) 21.5238095/21.5</p> <p>ECF for mp2 if height in cm² used</p> <p>Correct answer with no working gains 2 marks</p>	(2)

Question Number	Answer	Additional Guidance	Mark
2(c)(ii)	<p>An answer that makes reference to the following:</p> <ul style="list-style-type: none"> • females with Turner's syndrome have (slightly) lower mass and height (1) • but error bars overlap, so may be no difference/more data is needed (1) • cannot tell if BMI is significantly different as there are no error bars (1) • sample size is quite small (in both groups) (1) 		(4)

Question Number	Answer	Additional Guidance	Mark
3(a)(i)	<p>The only correct answer is A</p> <p>B is incorrect because bacteria have 70S ribosomes C is incorrect because bacteria do not have a nucleus D is incorrect because bacteria do not have a nucleus or 80S ribosomes</p>		(1)

Question Number	Answer	Additional Guidance	Mark
3(a)(ii)	<p>An answer that makes reference to the following:</p> <ul style="list-style-type: none"> • cytoplasm • plasmids • cell membrane 	<p>3 correct = 2 marks 1 or 2 correct = 1 mark</p> <p>ACCEPT glycogen granules / lipid droplets / mesosome</p>	(2)

Question Number	Answer	Additional Guidance	Mark
3(b)(i)	<ul style="list-style-type: none"> • 23 : 1.0 	Accept 23 : 1 Accept 1.0 : 0.043 Accept 1 : 0.043	(1)

Question Number	Answer	Additional Guidance	Mark
3(b)(ii)	An explanation that makes reference to the following: <ul style="list-style-type: none"> • <i>B. subtilis</i>, because it {is Gram positive / has a thick peptidoglycan cell wall} (1) • antibiotics interfere with the synthesis of peptidoglycan (crosslinks) / inhibit formation of peptidoglycan (1) 		(2)

Question Number	Answer	Additional Guidance	Mark
3(c)	An answer that makes reference to the following: <ul style="list-style-type: none"> • increase in cell number calculated (1) • rate calculated (1) 	$(1.7 \times 10^5 - 6 \times 10^3 =) 164\,000$ 1367 / 1366.7 / 1366.67 / 1366 recurring (cells per minute) 1366(.66) gains 1 mark Correct answer with no working gains 2 marks	(2)

Question Number	Answer	Additional Guidance	Mark
4(a)(i)	<p>The only correct answer is C</p> <p>A is incorrect because P is in prophase which comes before R which is in late anaphase</p> <p>B is incorrect because Q is in early anaphase which comes before R which is in late anaphase</p> <p>D is incorrect because S is in metaphase which comes before R which is in late anaphase</p>		(1)

Question Number	Answer	Additional Guidance	Mark
4(a)(ii)	<p>An answer that makes reference to the following:</p> <ul style="list-style-type: none"> • select the { anther / filament / carpel / ovary / embryo sac } (1) <p>and any four from:</p> <ul style="list-style-type: none"> • place in {acid / HCl} and {heat / place in hot / warm water bath} (1) • add acetic orcein (1) • tease the cells apart (with a mounted needle) / macerate (1) • place a coverslip on top and gently squash the preparation (1) • count total number of cells in each stage (of meiosis) / count (all) cells <u>and</u> count number of cells at metaphase I and metaphase II (1) • divide the number of cells in metaphase (multiplied by 100) by the total number of cells (1) 	<p>Sequence should be logical</p> <p>Accept stated temperature</p> <p>Accept other appropriate named stains e.g. Feulgen's, Toluene blue</p>	(5)

Question Number	Answer	Additional Guidance	Mark
4(b)	<p>An explanation that makes reference to the following:</p> <ul style="list-style-type: none"> • independent assortment (1) • crossing over between homologous chromosomes (1) • different combinations of <u>alleles</u> (in the gametes) / different combinations of maternal and paternal chromosomes (1) 		(3)

Question Number	Answer	Additional Guidance	Mark
5(a)	<p>An answer that makes reference to the following:</p> <p>A : matrix (space)</p> <p>B : outer membrane / envelope</p> <p>C : crista / cristae / (folded) inner membrane</p>	<p>Accept intermembrane space / double membrane</p> <p>All three correct = 2 marks 1 or 2 correct = 1 mark</p>	(2)

Question Number	Answer	Additional Guidance	Mark
5(b)	<p>An answer that makes reference to the following:</p> <ul style="list-style-type: none"> • diagram that (roughly) shows the shape and position of the cristae (minimum of 7 cristae at least half width of drawing) (1) • diagram that has no overlapping lines, breaks in lines, shading (1) 		(2)

Question Number	Answer	Additional Guidance	Mark
5(c)(i)	<p>The only correct answer is D</p> <p>A is incorrect because $74\,000 \div 0.9 = 82\,222$</p> <p>B is incorrect because $74\,000 \div 0.9 = 82\,222$</p> <p>C is incorrect because $74\,000 \div 0.9 = 82\,222$</p>		(1)

Question Number	Answer	Additional Guidance	Mark
5(c)(ii)	<p>An answer that makes reference to the following:</p> <ul style="list-style-type: none"> • 0.12 (195...) or 0.12 (162...) μm (1) 	<p>Accept ECF from Q5ci)</p> <p>A 8.33(3333) μm</p> <p>B 1.21(951...) or 1.21(621...) μm</p> <p>C 0.83(3333) μm</p>	(1)

Question Number	Answer	Additional Guidance	Mark
5(d)	<p>An explanation that makes reference to the following:</p> <ul style="list-style-type: none"> because (many) specimens / cells, will be colourless / not be visible (without a stain) (1) because electron microscope uses electrons to show (electron) dense areas (1) 	<p>Accept stains provide a contrast</p> <p>Accept electrons are not absorbed/scattered by coloured stains / electron microscopes need to use heavy metals</p>	(2)

Question Number	Answer	Additional Guidance	Mark
6(a)	<p>The only correct answer is A</p> <p>B is incorrect because Ebola has a helical capsid and an envelope whereas HIV has a polyhedral capsid</p> <p>C is incorrect because Ebola has a helical capsid and an envelope whereas λ (lambda) phage does not have an envelope and has a complex capsid</p> <p>D is incorrect because Ebola has a helical capsid and an envelope whereas TMV does not have an envelope</p>		(1)

Question Number	Answer	Additional Guidance	Mark
6(b)	<p>A description that makes reference to two of the following:</p> <ul style="list-style-type: none"> • globular / 3D / tertiary protein (1) • held together by {hydrogen bonds / disulfide bridges / ionic bonds/hydrophobic (interactions)} (1) • between R groups (1) • has an active site (1) 	<p>Accept hydrophobic R groups on inside /hydrophilic R groups on outside = mp 2 & 3</p>	<p>(3) EXP</p>

Question Number	Indicative content	
*6(c)	<p>Indicative content:</p> <p>Lytic Cycle (L)</p> <ul style="list-style-type: none"> • virus attaches/binds to (host) cells • (viral) genetic material / provirus, inserted (into host cell) • viral proteins/capsid made • genetic material / (viral) RNA, replicates • new viral particles assembled • cell ruptures/viral particles, released <p>Haemagglutinin (H)</p> <ul style="list-style-type: none"> • binds to sialic acid groups on the {glycoproteins / cell membrane / host cell / mucus} • so that it can <u>infect</u> host cell • needed by new virus particles to attach to more (host) cells <p>Neuraminidase (N)</p> <ul style="list-style-type: none"> • needed to separate virus particles from the {(non-host) cells / membrane} that they attach to • breaks down mucus • cleave the haemagglutinin from the sialic acid group on cell • needed to separate the new virus particles from each other • needed to separate the new virus particles from the host cell membrane (that they have budded out of) • needed to release the new virus particles from mucus • so that they are free to attach to new host cells • so that the infection is spread 	<p>Level 1:</p> <p>1 mark = 1 point from L, H or N</p> <p>2 marks = 2 points from L, H or N</p> <p>Level 2 :</p> <p>3 marks = 3 points from at least two of L, H or N</p> <p>4 marks = 4 points from at least two of L, H or N</p> <p>Level 3 :</p> <p>5 marks = 5 points from L, H and N</p> <p>6 marks = 6 points from L, H and N</p>

Question Number	Answer	Additional Guidance	Mark
7(a)	<p>An answer that makes reference to the following:</p> <ul style="list-style-type: none"> • a {vertical / upward sloping} line that shows the DNA content increasing (before mitosis) and a {vertical / downward sloping} line that shows the DNA content decreasing (after/towards end of mitosis) (1) • a horizontal line extending across all of mitosis (1) • DNA doubles after replication and returns to same level indicated at start (halves) (1) 		(3)

Question Number	Indicative content	
*7(b)	<p>Indicative content:</p> <p>DNA replication (D)</p> <ul style="list-style-type: none"> • reference to semi-conservative replication • helicase {separates the DNA strands / break hydrogen bonds between DNA strands} • each DNA strand acts as a template • bases complementary base pair with the exposed bases (on template strands) • DNA polymerase joins together (adjacent) DNA nucleotides • each new molecule (of DNA) has one old strand and one new strand • number of molecules doubles each time so total width of bands doubles each time <p>Stage 1 (S1)</p> <ul style="list-style-type: none"> • bases contain {heavy nitrogen / ^{15}N} • only one band near bottom of tube, as all the DNA has {heavy nitrogen / ^{15}N} <p>Stage 2 (S2)</p> <ul style="list-style-type: none"> • DNA replication occurs using the { light nitrogen / ^{14}N} • new strands will contain light nitrogen / ^{14}N} • only one band as all new molecules / DNA have one heavy and one light strand • band higher up in the tube as it is lighter than the {original / stage 1} DNA <p>Stage 3 (S3)</p> <ul style="list-style-type: none"> • new strands will contain { light nitrogen / ^{14}N} • (two bands because) {original / stage 1 / heavy nitrogen / ^{15}N strands} will bind with lighter strands to form {medium weight DNA molecules / band in same position as stage 2 } • (two bands because) light strands from stage 2 molecules will bind with new light strands to form lightest molecules, so will be higher up the tube • bands of equal widths as equal number of light and heavy strands in stage 2 DNA 	<p>Level 1: 1 mark = 2 details from D, S1, S2, S3 or S4</p> <p>2 marks = 3 details from D, S1, S2, S3 or S4</p> <p>Level 2: 3 marks = 4 details from two of D, S1, S2 S3 or S4</p> <p>4 marks = 4 details from three of D, S1, S2 S3 or S4</p> <p>Level 3: 5 marks = 5 details from four of D, S1, S2 S3 or S4</p> <p>6 marks = 5 details from all of D, S1, S2 S3 and S4</p>

	<p>Stage 4 (S4)</p> <ul style="list-style-type: none"> • all new strands contain { light nitrogen / ^{14}N } • (two bands because) {original / stage 1 / heavy nitrogen / ^{15}N strands} will bind with lighter strands to form {medium weight DNA molecules / band in same position as stage 2 / band in same position as the last stage} • (two bands because) light strands from stage 2 molecules will bind with new light strands to form lightest molecules, so will be { higher up the tube / in same position as the last stage } • more light strands in stage 3 DNA than heavy strands so the width of band for lightest DNA will be wider • in a ratio of 3 : 1 	
--	---	--

Question Number	Answer	Additional Guidance	Mark																								
8(a)	<table border="1" style="margin-left: auto; margin-right: auto;"> <thead> <tr> <th rowspan="2">Polysaccharide</th> <th colspan="4">Type of glycosidic bond</th> </tr> <tr> <th>both 1-4 and 1-6</th> <th>1-4 only</th> <th>1-6 only</th> <th>neither 1-4 nor 1-6</th> </tr> </thead> <tbody> <tr> <td>cellulose</td> <td style="text-align: center;"><input type="checkbox"/></td> <td style="text-align: center;">X</td> <td style="text-align: center;"><input type="checkbox"/></td> <td style="text-align: center;"><input type="checkbox"/></td> </tr> <tr> <td>glycogen</td> <td style="text-align: center;">X</td> <td style="text-align: center;"><input type="checkbox"/></td> <td style="text-align: center;"><input type="checkbox"/></td> <td style="text-align: center;"><input type="checkbox"/></td> </tr> <tr> <td>starch</td> <td style="text-align: center;">X</td> <td style="text-align: center;"><input type="checkbox"/></td> <td style="text-align: center;"><input type="checkbox"/></td> <td style="text-align: center;"><input type="checkbox"/></td> </tr> </tbody> </table>	Polysaccharide	Type of glycosidic bond				both 1-4 and 1-6	1-4 only	1-6 only	neither 1-4 nor 1-6	cellulose	<input type="checkbox"/>	X	<input type="checkbox"/>	<input type="checkbox"/>	glycogen	X	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	starch	X	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>		(3)
Polysaccharide	Type of glycosidic bond																										
	both 1-4 and 1-6	1-4 only	1-6 only	neither 1-4 nor 1-6																							
cellulose	<input type="checkbox"/>	X	<input type="checkbox"/>	<input type="checkbox"/>																							
glycogen	X	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>																							
starch	X	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>																							

Question Number	Answer	Additional Guidance	Mark
8(b)(i)	<p>A description that makes reference to the following:</p> <ul style="list-style-type: none"> deciding on colours is subjective / people may judge colours differently / hard to distinguish between some colours (1) the values for the concentration of sugars are ranges / semiquantitative / an { accurate / exact } sugar concentration can not be determined (1) 	<p>Accept idea that 1500 is both yellow and orange Accept idea that the ranges of concentration are large</p>	(2)

Question Number	Answer	Additional Guidance	Mark
8(b)(ii)	<p>An answer that makes reference to four of the following:</p> <ul style="list-style-type: none"> carry out test for reducing sugar (1) use chart to determine concentration of glucose / sucrose (1) carry out test for non-reducing sugar on the same volume of solution (1) heat in (HCl) acid (1) (then) neutralise (1) add same { volume/ number of drops } of Benedict's solution (1) subtract the concentration of glucose from the total concentration to determine sucrose concentration (1) 	<p>Accept description of test (Benedict's, plus heat -) for reducing and non-reducing</p> <p>Accept alternative method e.g mass or use of colorimeter</p> <p>Accept stated volumes</p> <p>Accept hydrolyse sucrose / sugar</p>	(4)

Question Number	Answer	Additional Guidance	Mark
9(a)(i)	<p>An explanation that makes reference to the following:</p> <ul style="list-style-type: none"> • phosphate heads are {polar / charged / hydrophilic}, so interact with / face the aqueous environment (inside and outside of cell) (1) • fatty acid tails form a {non-polar / hydrophobic } barrier / layer (on inside) (1) • (so control the) movement of { polar/charged / non-lipid soluble / water soluble } molecules into and out of the cell (1) 	<p>Accept waterproof layer</p> <p>Accept (only) allow lipid soluble molecules through</p>	(3)

Question Number	Answer	Additional Guidance	Mark
9(a)(ii)	<ul style="list-style-type: none"> <li data-bbox="421 308 1099 344">• number of phospholipids in $1 \mu\text{m}^2$ calculated (1) <li data-bbox="421 568 1055 604">• number of phospholipids in bilayer given (1) <li data-bbox="421 796 837 833">• answer in standard form (1) 	<p data-bbox="1240 272 1671 341">Example of calculation: $1\,000\,000 \div 0.3 = 3\,333\,333.3333$</p> <p data-bbox="1240 384 1821 414">1 mark for $1\,000\,000 \div 0.3 / 3\,333\,333.3333$</p> <p data-bbox="1240 564 1615 595">Accept x 2 as ecf from mp 1</p> <p data-bbox="1240 635 1868 665">$3\,333\,333.3333 \times 2 = 6\,666\,666.7 / 6\,666\,666.67$</p> <p data-bbox="1240 705 1715 735">2 marks for $6\,666\,667 / 6\,666\,666.67$</p> <p data-bbox="1227 783 1924 813">Accept answers in standard form as ecf for mp1 or 2</p> <p data-bbox="1240 853 1509 884">2 marks for 3.3×10^6</p> <p data-bbox="1240 924 1816 960">6.7×10^6 with no working out gains 3 marks</p>	(3)

Question Number	Answer	Additional Guidance	Mark
9(b)(i)	<p>A description that makes reference to the following:</p> <ul style="list-style-type: none"> • number of carbons/hydrogens in the fatty acid tail (1) • the number of carbon carbon double bonds (1) 	<p>Accept length of fatty acid tail /different R group chain</p> <p>Accept some are saturated / monounsaturated / polyunsaturated</p> <p>Accept different alcohol groups attached to phosphate</p> <p>Accept some have only one fatty acid tail and some have more</p>	(2)

Question Number	Answer	Additional Guidance	Mark
9(b)(ii)	<p>An answer that makes reference to three of the following:</p> <p>Similarities (max 2)</p> <ul style="list-style-type: none"> • all contain all four types of phospholipid (1) • A is the commonest in all four types of membrane (1) • B has the second highest percentage in all four types (1) <p>Differences (max 2)</p> <ul style="list-style-type: none"> • each membrane has a different { ratio/ percentage } of the four types of phospholipid (1) • C has the lowest percentage in the cell membrane and D has the lowest percentage in the other types of membrane (1) • the relative proportions of B, C and D are different in the cell membrane compared with the other three types of membrane (1) 	<p>Accept D is lowest in all except the cell membrane</p> <p>Accept description of the order of the types</p>	(3)

Question Number	Answer	Additional Guidance	Mark
9(b)(iii)	<p>An explanation that makes reference to the following:</p> <ul style="list-style-type: none">• when vesicles containing { proteins / (poly)peptides / lipids / steroids } from the endoplasmic reticulum fuse with the Golgi apparatus (1)• some of the membrane of the Golgi apparatus comes from the endoplasmic reticulum (1)	<p>Accept vesicles from ER move to the Golgi apparatus where proteins are modified</p>	<p>(2)</p>

